

Heart Rate during Group Flooding Therapy for PTSD

STEVEN H. WOODWARD, PH.D.,¹ KENT D. DRESCHER, PH.D.,¹ RONALD T. MURPHY, PH.D.,¹
JOSEPH I. RUZEK, PH.D.,¹ DAVID W. FOY, PH.D.,² NED J. ARSENAULT, B.A.,¹
FRED D. GUSMAN, L.C.S.W.¹

¹*National Center for PTSD, VA Palo Alto Health Care System, Palo Alto, CA*

²*Pepperdine University, Culver City, CA*

Abstract—The objective of this study was to assess, via heart rate, the arousal levels of participants in group trauma reexposure therapy for posttraumatic stress disorder, and so to better understand this common mode of treatment, particularly in regards to its presumed curative factor, extinction. Six Vietnam combat-related PTSD inpatients participated twice weekly in group trauma reexposure therapy during which their electrocardiograms were recorded. Heart rate was quantified continuously off-line. Heart rates of participants *not* directly engaged in imaginal reexposure to their personal combat traumas consistently exhibited mild linear declines from the beginnings to the ends of the approximately 2.5 hour sessions. Participants actively engaged in personal combat trauma reexposure exhibited higher whole-session heart rates. Most also exhibited more specific elevation extending over the later portions of therapy sessions during which intensive reexposure usually occurred. Surprisingly, no patients exhibited *focal* increases in heart rate concurrent with periods of most intensive traumatic incident review as judged from videotape. Administering imaginal reexposure in a group context does not preclude substantial physiological (sympathetic) arousal, as is preconditional for extinction. Under conditions in which the actively engaged reexposure patient is reliably identified, group trauma reexposure therapy may *not* provide an opportunity for “vicarious” flooding in nonengaged participants.

Key words—stress, traumatic, heart rate, group therapy

Introduction

AMONG THE TREATMENT APPROACHES to posttraumatic stress disorder (PTSD), controlled imaginal reexposure to the traumatic experience has arguably produced the most consistent benefits to patients (Blake et al., 1994; Solomon et al., 1992; Friedman and Southwick, 1995). Support for this approach has derived principally from clinical trials in which the treatment was delivered to individuals. Interest in the delivery of reexposure treatment to groups of traumatized patients has stemmed both from considerations of cost and availability, and from concern with patient retention and safety. Group treatment constitutes the primary modality by which trauma reexposure therapy is today administered by the largest provider of such services, the U.S. Department of Veterans Affairs. High drop-out rates have been observed from individual forms of trauma reexposure therapy (Foy et al., 1993), and serious complications have developed in some patients (Pitman et al., 1991). It has been proposed that group trauma reexposure therapy be explicitly constructed to incorporate the social support common to group treatment (Yalom, 1985) and a psychoeducational

Address for correspondence: Steven H. Woodward, Ph.D., NCPTSD Sleep Laboratory, VAMC 323-D6, 3801 Miranda Ave., Palo Alto, CA, 94304.

Integrative Physiological and Behavioral Science, January–March, 1997, Vol. 32, No. 1, 19–30.

component aimed at symptom management and relapse prevention. A large-scale clinical trial of group treatment for PTSD following these guidelines is now underway within the U.S. Department of Veterans Affairs (Cooperative Study #420).

An underlying assumption of reexposure therapies such as implosion and flooding is that reactivation of the trauma-related fear/anxiety state, with concomitant physiological (sympathetic) arousal, is a precondition for extinction to occur (Boudewyns and Shipley, 1983). We are aware of only one published study that actually recorded indices of sympathetic arousal *during* reexposure therapy with individual PTSD patients.¹ Meuser et al. (1991) reported on four cases of Vietnam combat-related PTSD whose heart rates were recorded during imaginal trauma reexposure. Their observations suggested that clear habituation of heart rate reactivity to imaginal stressors over sessions of reexposure treatment predicted therapeutic gain. Of particular interest here is whether the camaraderie and social support cultivated in group treatment could actually militate *against* the development of fear and anxiety states necessary for desensitization/extinction to occur. We tested this hypothesis by measuring heart rate (HR) during the course of manualized group trauma reexposure treatment delivered to Vietnam combat-related PTSD inpatients at the National Center for PTSD, Clinical Laboratory and Education Division.

Methods

Subjects

Eight consecutively-admitted male Vietnam combat-related PTSD inpatients were assigned to the treatment group via routine clinical decision-making. That is, all were expected to tolerate and derive benefit from trauma reexposure therapy. The assignment decisions were not based upon reactivity to trauma-related imagery either as measured or estimated from self-reports. The data reported here come from eleven of twenty-eight group treatment sessions and from six of the eight originally assigned patients. This sample of patients and sessions met the following criteria. Each patient participated in at least six group treatment sessions for which adequate ECG and video recordings were available and in which there could be identified a period of markedly intensive imaginal trauma reexposure as judged from videotape. Second, each patient, himself, underwent at least one episode of intensive imaginal trauma reexposure that was adequately recorded (in five of six cases, there were two such sessions). Of the original eight patients, one was eliminated because he had attended only four treatment sessions, and the other because his trauma review sessions were not videotaped. Of the remaining nineteen sessions, most did not include traumatic incident review as per the manualized treatment program. In sum, we present here data from eleven of sixteen (69 percent) programmed imaginal trauma reexposure episodes from this course of group trauma reexposure treatment. Thus, the data presented are likely to be representative of the sessions not recorded.

Of the six patients we report upon below, all met DSM-III-R criteria for PTSD assessed via the Clinician-Administered PTSD Scale (CAPS, Blake et al., 1991). As assessed via the Structured Clinical Interview for the DSM-III-R (SCID; Spitzer et al., 1990), all six also met criteria for recurrent major depressive disorder, only one being in remission at the time of his participation in this study. One patient met criteria for panic disorder with agoraphobia. He and one other patient met criteria for social phobia. Five of six patients exhibited a history of alcohol dependence, and four of six, of a non-alcohol substance dependence. These rates of comorbidity, though high, are fairly typical of chronic severe

PTSD patients (Kulka et al., 1988; McFarlane & Papay, 1992; Green et al., 1992; Engdahl et al., 1991). All six patients were taking psychotropic medications during the course of their participation in group trauma reexposure treatment. Four were medicated on trazodone, three on fluoxetine, two on buspirone, one on divalproex sodium, one on carbamazepine, and one on amitriptyline. One patient was taking two medications recognized to suppress heart rate reactivity (lisinopril 40 mg/day, nifedipine 60 mg/day).

Procedures

Subjects arrived at the treatment room five minutes prior to the scheduled start of the group. Subjects self-applied disposable silver/silver-chloride electrodes to the medial surfaces of each wrist, chose a seat in the room, and connected their electrodes to leads hanging behind each chair. Though wrist derivations were not optimal from the standpoint of artifact, we assigned a high priority to minimizing the intrusiveness of our recording methods. The electrode leads were routed around the outside of the room to an electrode junction box mounted on the wall. A heavily shielded cable then conducted the ECG signals to a Grass Model 78 polygraph in the next room. ECG was filtered to a 10–30 Hz bandwidth prior to digitization. Again, though not ideal, these filtering parameters allowed optimal resolution of R-waves under the difficult recording conditions involved. The ECG signals were digitized at 125 Hz and streamed to disk using a data-acquisition interface to a laboratory computer (DATAQ Instruments, Akron, OH). A computer-based algorithm calculated R-R intervals and HRs on a four-second time-base. The use of a short time-base allowed us to more easily exclude artifactual epochs. This was accomplished by first requiring that the standard deviation of the interbeat interval be less than 10 msec. As both 60 Hz noise and high-frequency high-amplitude muscle artifact could produce signal epochs passing this criterion, a combination of range-checking and review of raw data was applied to the remaining epochs. Linear interpolation between nonartifactual epochs followed by a 15-point “boxcar” moving average filter yielded the final HR time-series. Both digital ECG records and videotape were time-stamped, and their registration was maintained within one second.

Group Trauma Reexposure Therapy Protocol

Patients at the National Center for PTSD, Clinical Laboratory and Education Division, are involved in an intensive inpatient program that emphasizes psychoeducational and process-oriented group psychotherapy aimed at reducing PTSD symptoms and treating comorbid disorders such as depression and substance abuse. At the time of this study, patients typically resided in the program for six months. After two to three months, patients were assigned to a “Trauma Focus” group. According to the manualized protocol administered to the patients in the current study, the initial set of group sessions were psychoeducational in format, intended to provide a cognitive structure incorporating PTSD, combat trauma, and coping, that supported the coming reexposure phase. The reexposure phase of treatment was then accomplished one patient at a time. Therapists guided each individual through an imaginal reexperience of his most severe combat traumas in such a way as to elicit and maintain the expression of distressing affects associated with those events. Two sessions were devoted to each individual’s combat traumas. At the end of each reexposure session, the nonengaged participants were debriefed regarding the thoughts and emotions stimulated by the activity. The reexposure phase of treatment was

followed by an integrative phase focusing upon relapse prevention and the maintenance of therapeutic gains.

Results

Mean Combat Exposure Score (CES; Keane et al., 1989) for the sample was 24.8 (s.d. 8). Mean score on the Mississippi Scale for Combat-Related PTSD (MISS) was 133.8 (s.d. 15.4). Beck Depression Inventory (BDI, Beck et al., 1961) scores were, on average, 29.8 (s.d. 9.3). These indices and the levels of comorbidity discussed above confirm that the participants in this study were highly symptomatic.

Figure 1 presents HR profiles for each of the eleven sessions analyzed. For each session, the HR of the individual undergoing trauma reexposure is plotted (in black) separately from the mean of the other participants (in grey). Also indicated via (grey) vertical lines are the break periods which typically occurred approximately one third of the way through the sessions, and (in black) the episodes of intensive imaginal re-exposure as judged from videotape. In nine of the eleven sessions reported upon here, these episodes occurred during the second two-thirds of the session. Clock time is represented along the x-axis. It can be seen that most sessions lasted approximately 2.5 hours.

The mean HR profiles of the participants not undergoing imaginal trauma reexposure were markedly consistent in exhibiting mild near-linear declines from the beginning to the ends of sessions. These declines typically amounted to approximately 5 beats per minute (BPM). (Variation in the mean levels of profiles in Figure 1 derives from the fact that different subgroups of patients contributed from session to session.) The continuity of these declines across break periods was notable in light of the frequency with which participants smoked cigarettes during breaks. The HR profiles of individuals undergoing imaginal reexposure were highly variable: some exhibited large increases in the later portions of treatment sessions during which most intense imaginal reexposure occurred; others exhibited relatively flat profiles. Consistent across these profiles, however, was the absence of any focused excursion in HR coincident with the period of most intense imaginal reexposure. Figure 2 portrays this point more strongly by plotting the mean of all eleven HR profiles recorded from participants undergoing imaginal reexposure *centered* on the midpoints of those periods judged most intense from videotape. As the most intense periods were typically only a few minutes long, the 26-minute epoch plotted in Figure 2 would seem to provide ample time to visualize any phasic evolution and resolution of arousal coincident with most intensive reexposure. No such morphology was in evidence. Figure 2 also reinforces a second feature of the individual plots in Figure 1, that the HR time series of group members not on the "hot seat" exhibited no reexposure-related morphology whatsoever.

Figure 3 presents the same HR data in another way, contrasting each individual's mean HR profile from sessions in which he was the actively engaged reexposure patient with his mean profile from sessions in which he was not. (As session lengths differed somewhat, these plots are truncated to the length of the shorter of the two therapy sessions in which the patient was the actively engaged patient, that is, for the five of six patients who provided two such sessions. As well, the terminal sections of nonreexposure session mean profiles necessarily represent averages taken over variable numbers of sessions.) Appreciable in Figure 3 is the overall consistency with which group participants' HRs were elevated when they were identified as the patient engaged in imaginal reexposure as compared to when they were not so identified. In four of six patients, this elevation






Fig. 1. All eleven group trauma reexposure sessions are represented in this figure. Each subplot contains two HR time series. The grey time series is the average computed over all patients not actively engaged in trauma reexposure during that session. The black time series plots the HR of the single patient who was so engaged. The black vertical lines indicate the period of intense imaginal reexposure judged from videotape. The grey vertical lines indicate the break period. Clock time is noted below the x-axis. Here, as in all subsequent figures, the y-axis represents HR in BPM.


Session 11/29/95



Session 12/06/95



Session 11/29/95



Session 12/05/95




Fig. 1. (Continued)




Fig. 1. (Continued)




FIG. 2. Average of all HR profiles in Fig. 1 centered on periods of most intense imaginal reexposure as judged from videotape. The grey time series is the average over all sessions in which patients were not actively engaged in trauma reexposure. The black time series is the average over the session(s) in which the patients were so engaged. Total epoch duration is 26 minutes. Note the absence of any peaking in HR associated with periods of most intense imaginal reexposure.

favoring the latter two-thirds of sessions during which most intense reexposure generally occurred. A fifth patient exhibited an elevation that was not specific to the latter two-thirds of sessions, while the sixth showed very little HR variation of any kind. It was later determined that this last patient was taking both an antihypertensive and an antianginal medication.

Figure 4 portrays the HR data averaged by the first, second, and third third of each session. A univariate repeated-measures ANOVA indicated that the interaction of session type (reexposure vs. nonreexposure) with time (first, second, third) was statistically significant ($F(2,10) = 23.99$, $p < 0.025$, $\epsilon = 0.7002^2$). The main effects of time ($F(2,10) = 8.95$, $p < 0.029$) and session type ($F(1,5) = 6.39$, $p < 0.053$) were also statistically significant or nearly so despite the small sample size. Computing univariate contrasts, it was determined




FIG. 3. Each of the above plots contains two averaged HR time series from a single patient. The grey time series is the average over all sessions in which the patient was not actively engaged in trauma reexposure. The black time series is the average over the session(s) in which the patient was so engaged. Note the tendency for HR to be elevated in the latter portions of sessions, during which most intensive imaginal reexposure occurred. Note also the variation across patients in exposure-driven HR, as contrasted with the consistent mild decline of HR during nonengaged sessions.




FIG. 4. Mean HR averaged over subjects and sessions, separated by session thirds. Grey bars represent the nonactively engaged condition, black bars, the actively engaged condition.

that the session type effect was significant at the second ($F(1,5) = 8.94$, $p < 0.03$) and third ($F(1,5) = 8.43$, $p < 0.034$) third of sessions, but not at the first ($F(1,5) = 1.54$, n.s.). Considering only nonreexposure HR, the reduction from the first third to the second was significant ($F(1,5) = 21.79$, $p < 0.005$) and the reduction from the second third to the third, nearly so ($F(1,5) = 5.10$, $p < 0.07$). Considering HR during reexposure sessions, the increase from the first third to the second was significant ($F(1,5) = 8.56$, $p < 0.033$), while the other contrasts were not (2 vs. 3: $F(1,5) = 0.603$, n.s.; 1 vs. 3: $F(1,5) = 0.601$, n.s.).

We also analyzed the HR data for their capacity to predict short-term benefit from group trauma reexposure therapy. Of particular interest to us was whether reactivity indicated by HR elevation during the latter portions of sessions was associated with benefit from reexposure treatment. Figure 3 is ordered according to the subjective salience of the HR reactivity following this pattern. While more objective and elaborate methods could have been used to quantify the magnitude of "slow" curvature in the HR time series, the small number of patients involved did not justify their application here. (Any such quantitative index would be retransformed to a simple ranking for subsequent analyses.) Two of us (RTM and JIR) who served as the group therapists in the study ranked the patients according to apparent global benefit from trauma reexposure. These rankings were performed independently and blind to the HR data. We also examined MISS scores acquired from patients before and after group treatment.

Remarkably close agreement was found between therapists' rankings of global benefit from trauma reexposure and independent rankings of apparent treatment-driven reactivity in individuals' HR profiles (Fig. 3; Spearman's $\rho = 0.943$, $p < 0.01$). After debriefing the therapists (RTM and JIR), it became evident that a patient's display of overt congruent

affect during imaginal trauma reexposure was the major factor underlying their rankings. Thus, perceived affectivity during reexposure corresponded to an index of HR reactivity. An examination of MISS scores pre- and post-group uncovered no direct correspondence between our ranking of HR reactivity and treatment benefit as indexed by self-reported PTSD symptomology. In fact, the relation between these rankings was negative (Spearman's $\rho = -0.70$, n.s.). That is, the patient who had the most HR reactivity during imaginal reexposure (and the most overt congruent affectivity according to the group therapists) actually reported worsening in PTSD symptom severity on the MISS. Conversely, the patient with the least HR reactivity reported the most improvement in PTSD symptom severity.

Discussion

This study concerns the performance of only six patients engaged in group trauma reexposure therapy, so its conclusions must be regarded as provisional in that light. Acknowledging that caveat, important aspects of the data derived from this group appeared unambiguous. Substantial sympathetic arousal was observed in the group therapy context in those individuals actively engaged in trauma reexposure treatment. Participants' whole-session HRs were 3 BPM higher when they were the actively engaged reexposure patient than when they were not. The average differential rose to 8 BPM during the latter portions of sessions during which most intensive reexposure occurred. Not only were these differences during the latter portions of groups statistically significant, but they were larger than the HR increases typically observed in association with trauma cues in reactivity studies (Prins et al., 1995). In some cases, they were also sustained over substantial periods of time. By the same token, the differences in HR between engaged and nonengaged conditions was equally the result of the absence of HR elevation during the latter. Group members not actively engaged by the therapist in personal trauma reexposure demonstrated a gradual reduction in arousal as indexed by HR that extended throughout the sessions, including the brief episodes of most intense reexposure and traumatic incident review. Clearly, "vicarious flooding" was not evidenced by these six patients. In light of the fact that the nonengaged patients were in close proximity to and listening to a peer describing traumatic experiences similar to their own, the contrast between theirs' and the HR profiles of engaged patients is impressive. Assuming these six patients are not unique, a better understanding of the "mental set" underlying and sustaining *nonarousal* in some participants of trauma reexposure group therapy may be necessary to optimize this form of treatment.

A modest examination of relationships between HR reactivity during trauma reexposure and therapeutic gain produced a number of suggestive findings, which again must be considered in light of the small size of the study sample. When therapists were asked to estimate benefit from reexposure, their judgments corresponded remarkably to a ranking of "exposure-driven" HR reactivity derived from the HR time series (Fig. 3). On debriefing, it was determined that the therapists had spontaneously transformed "benefit from exposure" to "affectivity during exposure." In light of the broadly-held model of exposure therapy that regards physiologic (sympathetic) arousal as a precondition for extinction, the apparent accuracy with which two therapists could assess arousal via affectivity is heartening. Unfortunately, there were no such relationships observed between the rankings of arousal and affectivity and self-reported change in PTSD severity pre- and post-treatment. In fact, a trend toward the reverse relationship was observed. At least two possibilities should be

considered in regards to this finding. It may be that those patients most "successfully" engaged in reexposure treatment will experience a temporary overwhelming of avoidant coping responses with temporary increases in PTSD symptom severities, but greater long-term therapeutic gain. This possibility deserves close scrutiny in future outcome research, particularly in light of the findings of Pitman et al. (1991), as it implies that the patients who "perform best" in reexposure treatment may be precisely those requiring the most intense post-exposure treatment follow-up. A second, more disturbing possibility is that, given the apparent absence of "vicarious flooding," the participants in this course of group therapy received insufficient numbers of reexposure episodes to promote extinction. Instead, individuals experienced sensitization in approximate proportion to the levels of physiologic reactivity and affectivity induced by reexposure. This possibility also deserves close scrutiny in future outcome research.

Finally, important improvements could be made with relative ease to the methodology we used in this study. First, this study relied almost exclusively on physiologic reactivity as a process measure, whereas both subjective distress and behavioral avoidance have been shown to be important parallel indices of reactivity during exposure therapy. Second, informal inquiries of patients made during the course of this study suggested that standard chest leads could have been used without undue intrusion upon the group process. In fact, the patients in this group interpreted the physiological recording not as intrusive but, rather, as indicative of the seriousness of the therapeutic enterprise in which they were engaged. Standard disposable ECG electrodes can be applied with little skin preparation and will produce stronger and less artifact-laden ECG signals than we obtained. Second, small sensitive lapel microphones could be placed so as to record breath sounds and speech. This signal could then be integrated and digitized at rates comparable to the EEG in order to provide both respiratory rate and speech rate information. While a high speech rate was noted in only one of the patients of this study (who did not stand as an extreme scorer either with regard to baseline HR or HR reactivity), the continuous indexing of speech rate as a contributor to cardiac rate elevation could be useful. The indexing of respiratory rate could provide important utility in that its combination with high-quality ECG signal would enable estimation of the respiratory sinus arrhythmia (Grossman et al., 1991), a continuous index of parasympathetic drive on the heart. By independently estimating parasympathetic drive, a richer picture of arousal status during trauma reexposure treatment could be derived from the ECG (Berntson et al., 1991, 1993; Porges, 1995).

Acknowledgment

The authors wish to acknowledge the Research Service of the Veterans Affairs Palo Alto Health Care System for administrative support of this pilot project.

Notes

1. One such study is in progress (Resick, personal communication).
2. Huynh-Felt corrections for nonsphericity used.

References

- American Psychiatric Association (1952). *Diagnostic and Statistical Manual of Mental Disorders*. Washington: American Psychiatric Association.
- American Psychiatric Association (1968). *Diagnostic and Statistical Manual of Mental Disorders*. Washington: American Psychiatric Association. Ed. 2.
- American Psychiatric Association (1980). *Diagnostic and Statistical Manual of Mental Disorders*. Washington: American Psychiatric Association. Ed. 3.
- American Psychiatric Association, (1987). *Diagnostic and Statistical Manual of Mental Disorders*. Washington, DC: American Psychiatric Association. Ed. 3r.
- American Psychiatric Association (1994). *Diagnostic and Statistical Manual of Mental Disorders*. Washington, DC. Ed. 4.
- Achenbach, T.M. & Edelbrock, C.S. (1983). *Manual for the Child Behavior Checklist and Revised Child Behavior Profile*. New York: Queen City Printers.
- Achenbach, T.M. (1991). *Manual for the Child Behavior Checklist/4-18 and 1991 Profile*. Burlington, VT: University of Vermont, Dept. of Psychiatry.
- Ackerman, P.T., Dykman, R.A., Holcomb, P.J., & McCray, D.S. (1982). Methylphenidate effects on cognitive style and reaction time in four groups of children. *Psychiatry Research*, 7, 199-213.
- Angel, C., DeLuca, D.C., Newton, J.E.O., & Reese, W.G. (1982). Assessment of pointer dog behavior: Drug effects and neurochemical correlates. *Pavlovian Journal of Biological Science*, 17.
- Antelman, S. (1988). Stressor-induced sensitization to subsequent stress: Implications for the development and treatment of clinical disorders. In P.W. Kalivas and C.D. Barnes (Eds.), *Sensitization the Nervous System*. Caldwell, NJ: Telford Press, pp. 227-257.
- Attias, J., Bleich, A., Furman, V., & Zinger, Y. (1996). Event-related potentials in post-traumatic stress disorder of combat origin. *Biological Psychiatry*, 40:373-381.
- Attias, J., Bleich, A., & Gilat, S. (1996). Classification of veterans with post-traumatic stress disorder using visual brain evoked P3s to traumatic stimuli. *British Journal of Psychiatry*, 168:100-115.
- Beck, A.T.; Ward, C.H.; Mendelson, M.; Mock, J.; Erbaugh, J. (1961) An inventory for measuring depression. *Archives of General Psychiatry*, 4:561.
- Becker, J.V., Skinner, L.J., Abel, G.G., Axelrod, R., & Cichon, J. (1984). Sexual problems of sexual assault survivors. *Women and Health*, 9, 5-20.
- Begleiter, H. & Platz, A. (1969). Cortical evoked potentials to semantic stimuli. *Psychophysiology*, 6:91-100.
- Berntson, G.G., Cacioppo, J.T., Quigley, K.S. (1991). Autonomic determinism: The modes of autonomic control, doctrine of autonomic space, and the laws of autonomic constraint. *Psychological Review*, 98:459-487.
- Berntson, G.G., Cacioppo, J.T., Quigley, K.S. (1993). Respiratory sinus arrhythmia: Autonomic origins, physiological mechanisms, and psychophysiological implications. *Psychophysiology*, 30:183-196.
- Blake, D.D., Weathers, F., Nagy, L.M., Kaloupek, D.G., Klauminizer, G., Charney, D.S., Keane, T.M. (1991). A clinician rating scale for assessing current and lifetime PTSD: The CAPS-1. *Behavior Therapist*, 14:187-188.
- Blake, D.D., Abueg, F.R., Woodward, S.H. (1994). Treatment efficacy in posttraumatic stress disorder. In T.R. Giles (Ed.) *Handbook of Effective Psychotherapy*. New York: Plenum.
- Blanchard, E.B., Kolb, L.C., Pallmeyer, T.P., & Gerardi, R.J. (1982). A psychophysiological study of posttraumatic stress disorder in Vietnam veterans. *Psychiatric Quarterly*, 54, 220-229.
- Blanchard, E.B., Kolb, L.C., Gerardi, R.J., Ryan, P., & Pallmeyer, T.P. (1986). Cardiac response to relevant stimuli as an adjunctive tool for diagnosing post-traumatic stress disorder in Vietnam veterans. *Behavior Therapy*, 17, 592-606.
- Blanchard, E.B., Kolb, L.C., Prins, A. (1991). Psychophysiological responses in the diagnosis of posttraumatic stress disorder. *Journal of Nervous and Mental Disorders*, 179:97-101.
- Boudewyns, P., Shipley, R.H. (1983). *Flooding and implosive therapy*. New York: Plenum.
- Bremner, J.D., Scott, T.M., Delaney, R.C., Southwick, S.M., Mason, J.W., Johnson, D.R., Innis, R.B., McCarthy, G., Charney, D.S. (1993). Deficits in Short-term Memory in Posttraumatic Stress Disorder. *American Journal of Psychiatry*, 150:1015-1019.

- Bremner, J.D., Southwick, S.M., Johnson, D.R., Yehuda, R., & Charney, D.S. (1993). Childhood physical abuse and combat-related posttraumatic stress disorder in Vietnam veterans. *American Journal of Psychiatry*, 150, 235-239.
- Breslau, N., Davis, G.C., Andreski, P., & Peterson, E. (1991). Traumatic events and posttraumatic stress disorder in an urban population of young adults. *Archives of General Psychiatry*, 48, 216-222.
- Brett, E.A. (1993). Classifications of posttraumatic stress disorder in DSM-IV: Anxiety disorder, dissociative disorder, or stress disorder? In J.R.T. Davidson and E.B. Foa (Eds.), *Posttraumatic Stress Disorder. DSM-IV and Beyond*. Washington: American Psychiatric Press, p. 191-206.
- Briere, J.N. & Elliott, D.M. (1994). Immediate and long-term impacts of child sexual abuse. In R.E. Behrman (Ed.), *The Future of Children*. Los Altos, CA: Center for the Future of Children, The David and Lucile Packard Foundation, pp. 54-69.
- Browne, A. & Finkelhor, D. (1986) Impact of child sexual abuse: A review of the research. *Psychological Bulletin*, 99, 66-77.
- Bryant, R.A. & Harvey, A.G. (1995). Processing threatening information in posttraumatic stress disorder. *Journal of Abnormal Psychology*, 104:537-541.
- Buchsbaum, M., & Silverman, J. (1968). Stimulus intensity control and the cortical evoked response. *Psychosomatic Medicine*, 30:12-22.
- Cassiday, K.L., McNally, R.J., & Zeitlin, S.B. (1992). Cognitive processing of trauma cues in rape victims with posttraumatic stress disorder. *Cognitive Therapy and Research*, 16:283-295.
- Charles, G., Hansenne, M., Ansseau, M., Pitchot, W., Machowski, R., Schittecatte, M., & Wilmotte, J. (1995). P300 in posttraumatic stress disorder. *Neuropsychobiology*, 32:72-74.
- Charney, D.S., Deutch, A.Y., Krystal, J.H., Southwick, S.M., & Davis, M. (1993). Psychobiologic mechanisms of posttraumatic stress disorder. *Archives of General Psychiatry*, 50, 294-305.
- Cicchetti, D. & Toth, S.L. (1995). A developmental psychopathology perspective on child abuse and neglect. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34, 541-565.
- Coles, M.G.H., Gratton, G., Bashore, T.R., Eriksen, C.W., & Donchin, E. (1985). A psychophysiological investigation of the continuous flow model of human information processing. *Journal of Experimental Psychology: Human Perception and Performance*, 11:529-553.
- Conte, J. & Schuerman, J. (1987). Factors associated with an increased impact of child sexual abuse. *Child Abuse and Neglect*, 11:201-211.
- Cornblatt B.A., Lenzenweger M.F., Erlenmeyer-Kimling L. (1989). The Continuous Performance Test, Identical Pairs Version: II. Contrasting attentional profiles in schizophrenic and depressed patients. *Psychiatry Research*, 29:65-85.
- Cornblatt B.A., Kelip J.C. (1994). Impaired attention, genetics, and the pathophysiology of schizophrenia. *Schizophrenia Bulletin*, 20(1):31-46.
- D'Amato, R.J., Blue, M.E., Largent, B.L. (1987). Ontogeny of the serotonergic projection to rat neocortex: Transient expression of a dense innervation to primary sensory areas. *Proceedings of the National Academy of Science*, 84, 4322-4326.
- Dalton J.E., Pederson S.L., Blom B.E., Besyner J. (1986). Neuropsychological screening for Vietnam veterans with PTSD. *VA Practitioner*, 3(7):37-47.
- Davidson, J.R.T. (1993). Issues in the diagnosis of Posttraumatic stress disorder. In J. M. Oldham, M. R. Riba, and A. Tasman (Eds.), *Review of Psychiatry*. Washington: American Psychiatric Press, pp. 141-156.
- Davidson, J.R.T. & Foa, E.B. (1993). *Posttraumatic Stress Disorder. DSM-IV and Beyond*. Washington: American Psychiatric Press.
- DeBellis, M.D., Lefter, L., Trickett, P.K., & Putnam, F.W. (1994). Urinary catecholamine excretion in sexually abused girls. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33, 320-327.
- Deblinger, E., McLeer, S.V., Atkins, M.S., Ralphe, D., Foa, E. (1989). Post-traumatic stress in sexually abused, physically abused, and non-abused children. *Child Abuse and Neglect*, 13, 403-408.
- Donchin, E. & Coles, M. (1988). Is the P300 component a manifestation of context updating? *Behavioral and Brain Sciences*, 11:357-374.
- Duncan-Johnson, C. C., & Kopell, B. S. (1981). The Stroop effect: Brain potentials localize the source of interference. *Science*, 214:938-940.
- Dykman, R.A. & Gantt, W.H. (1951a). A comparative study of cardiac conditioned responses and motor conditioned responses in controlled "stress" situations. *American Psychologist*, 6:263.
- Dykman, R.A. & Gantt, W.H. (1951b). A comparative study of cardiac and motor conditional responses. *American Journal of Physiology*, 167.
- Dykman, R.A. & Gantt, W.H. (1954). Blood pressure conditioned to pain. *Federal Proceedings*.

- Dykman, R.A., Gantt, H., & Whitehorn, J.C. (1956). Conditioning as emotional sensitization and differentiation. *Psychology Monographs*, 7:1-17.
- Dykman, R.A. & Gantt, W.H. (1960). A case of experimental neurosis and recovery in relation to the orienting response. *Journal of Psychology*, 50:105-110.
- Dykman, R.A., Corson, S.A., Reese, W.G., & Seager, L.D. (1962). Inhibition of urine flow as a component of the conditioned defense reaction. *Psychosomatic Medicine*, 24:177-186.
- Dykman, R.A., Murphree, O.D., & Peters, J.E. (1969). Like begets like: Behavioral tests, classical autonomic and motor conditioning, and operant conditioning in two strains of pointer dogs. *Annals of the New York Academy of Sciences*, 159:976-1007.
- Dykman, R.A., Ackerman, P.T., Holcomb, P.J., & Boudreau, A.Y. (1983a). Physiological manifestations of learning disability. *Journal of Learning Disabilities*, 16:46-53.
- Dykman, R.A., Holcomb, P.J., Ackerman, P.T., & McCray, D.S. (1983b). Auditory ERP augmentation-reduction and methylphenidate dosage needs in attention and reading disabled children. *Psychiatry Research*, 9:255-269.
- Dykman, R.A., Ackerman, P.T., and Newton, J.E.O. (1996). Post-traumatic stress disorder: A sensitization reaction. *Integrative Physiological and Behavioral Sciences*, (this volume).
- Engdahl, B.E., Speed, N., Eberly, R.E., Schwartz, J. (1991). Comorbidity of psychiatric disorders and personality profiles of American World War II prisoners of war. *Journal of Nervous and Mental Disease*, 1179:181-87.
- Epstein, A.W. (1964). Recurrent Dreams. *Archives of General Psychiatry*, 10:25-30.
- Epstein, A.W. and Hill, W. (1966). Ictal phenomena during REM sleep of a temporal lobe epileptic. *Archives of Neurology*, 15:367-375.
- Eriksen, B.A., & Eriksen, C.W. (1974). Effects of noise letters upon the identification of target letter in a nonsearch task. *Perception and Psychophysics*, 16:143-149.
- Everly, G.S. (1993). Psychotraumatology: A two factor formulation of posttraumatic stress. *Integrative Physiological and Behavioral Science*, 28:270-278.
- Eysenck, H.J. (1979). The conditioning model of neurosis. *Behavioral and Brain Sciences*, 2:155-199.
- Fairbank, J., Keane, T. and Malloy, P. (1983). Some preliminary data on the psychological characteristics of Vietnam veterans with posttraumatic stress disorders. *Journal of Consulting and Clinical Psychology*, 6:912-919.
- Fairbank, J., McCaffrey, R. and Keane, T. (1985). Psychometric detection of fabricated symptoms of posttraumatic stress disorder. *American Journal of Psychiatry*, 142:501-503.
- Fairbank, J.A. & Brown, T.A. (1987). Current behavioral approaches to the treatment of posttraumatic stress disorder. *Behavior Therapist*, 10:57-64.
- Famularo, R., Kinscherff, R., Fenton, T. (1992). Psychiatric diagnoses of maltreated children: Preliminary findings. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31:863-867.
- Famularo, R. & Fenton, T. (1994). Early development history and pediatric posttraumatic stress disorder. *Archives of Pediatric and Adolescent Medicine*, 148:1032-1038.
- Faux, S.F., Shenton, M.E., McCarley, R.W., Torello, M.W. and Duffy, F.H. (1987). P200 topographic alterations in schizophrenia: Evidence for left temporal-centroparietal region deficits. In R. Johnson R, J.W. Rohrbaugh, and R. Parasuraman (Eds.). *Current Trends in Event-Related Potential Research (EEG Suppl. 40)*. New York: Elsevier, pp. 681-687.
- Figley, C.R. (1986) *Trauma and its wake: Traumatic stress theory, research, and intervention*. New York, NY: Brunner/Mazel.
- Foa, E.B. & Kozak, M.J. (1986). Emotional processing of fear: Exposure to corrective information. *Psychological Bulletin*, 99:20-35.
- Foa, E.B., Steketee, G., & Rothbaum, B.O. (1989). Behavioral/cognitive conceptualizations of post-traumatic stress disorder. *Behavior Therapy*, 20:155-176.
- Foa, E.B., Feske, U., Murdock, T.B., Kozak, M.J., McCarthy, P.R. (1991). Processing of threat-related information in rape victims. *Journal of Abnormal Psychology*, 100(2):156-162.
- Foa, E.B. & Riggs, D.S. (1995). Posttraumatic stress disorder following assault: Theoretical considerations and empirical findings. *Current Directions*, 4:61-65.
- Foy, D.W., Kaga, R., Haas, B., Leski, G., Hammen, M., Smith, R., Reno, S. (1993). Parameters of flooding therapy with chronic combat-related PTSD: High vs. Low dose comparisons. Paper presented to the International Society for Traumatic Stress Studies, San Antonio, TX.

- Friedman, M.J. & Southwick, S. (1995). Towards pharmacotherapy for posttraumatic stress disorder. In M.J. Friedman, D.S. Charney, A. Deutch (Eds.) *Neurobiological and Clinical Consequences of Stress: From Normal Adaptation to PTSD*. Philadelphia: Lippincott-Raven Publishers, pp. 465-482.
- Gantt, W.H. (1944). *Experimental Basis for Neurotic Behavior*. New York: Hoeber.
- Gantt, W.H. & Dykman, R.A. (1952). Experimental psychogenic tachycardia. *American Journal of Physiology*, 171.
- Gantt, W.H. (1953). Principles of nervous breakdown-schizokinesis and autokinesis. *Annals of New York Academy of Sciences*, 56:143-163.
- Gil, T., Calec, A., Greenberg, D. (1990). Cognitive functioning in posttraumatic stress disorder. *Journal of Traumatic Stress*, 1:29-45.
- Goddard, G., McIntyre, D., and Leech C. (1969). A permanent change in brain function resulting from daily electrical stimulation. *Experimental Neurology*, 25:295-330.
- Golier, J., Yehuda, R., Cornblatt, B., Harvey, P., Gerber, D., and Levengood, R. (1996). Sustained attention in combat-related posttraumatic stress disorder. *Integrative Physiological and Behavioral Sciences* (this volume).
- Gray, J.A. (1982a). *The Neuropsychology of anxiety: An inquiry into the function of the septo-hippocampal system*. Oxford: Oxford University Press.
- Gray, J.A. (1982b). Precis of the Neuropsychology of Anxiety: An Inquiry into the Functions of the Septo-hippocampal System. *Behavioral and Brain Sciences*, 5:469-534.
- Green, A.H. (1993). Child sexual abuse: Immediate and long-term effects and intervention. *Journal of the American Academy of Child and Adolescent Psychiatry*, 32:890-902.
- Green, B.L., Lindy, J.D., Grace, M.C., Leonard, A.C. (1992). Chronic posttraumatic stress disorder and diagnostic comorbidity in a disaster sample. *Journal of Nervous and Mental Disease*, 180:760-6.
- Greenhouse, S.W. & Geisser, S. (1959). On methods in the analysis of profile data. *Psychometrika*, 24:95-112.
- Grossman, P., Karemaker, J., Wieling, W. (1991). Prediction of tonic parasympathetic cardiac control using respiratory sinus arrhythmia: The need for respiratory control. *Psychophysiology*, 28:201-216.
- Gurvits, T.V., Lasko, N.B., Schachter, S.C., Kuhne, A.A., Orr, S.P., Pitman, R.K. (1993). Neurological status of Vietnam veterans with chronic Posttraumatic Stress Disorder. *Journal of Neuropsychiatry and Clinical Neurosciences*, 5:183-88.
- Hamilton, J. (1985). Pseudo-Posttraumatic Stress Disorder. *Military Medicine*, 150:353-356.
- Hebb, D.O. (1949). *The Organization of Behavior. A Neuropsychological Theory*. New York:Wiley.
- Hegerl, U. & Juckel, G. (1993). Intensity dependence of auditory evoked potentials as an indicator of central serotonergic neurotransmission: A new hypothesis. *Biological Psychiatry*, 33:173-187.
- Herman, J.L. (1993). Sequelae of Prolonged and Repeated Trauma: Evidence for a Complex Posttraumatic Syndrome (DESNOS). In J.R.T. Davidson and E.B. Foa (Eds.). *Posttraumatic Stress Disorder: DSM-IV and Beyond*. Washington: American Psychiatric Press, pp. 213-228.
- Herman-Giddens, M.E., Sandler, A.D., & Friedman, N.E. (1988). Sexual precocity in girls: An association with sexual abuse? *American Journal of Diseases of Children*, 142:431-433.
- Heur, F., & Reisberg, D. (1992). Emotion, arousal, and memory for detail. In S. Christianson (Ed.). *The handbook of emotion and memory: Research and theory*. Hillsdale, NJ: Erlbaum, pp. 151-80.
- Hilgard, E.R. & Bower, G.H. (1975). *Theories of Learning*. Englewood Cliffs, NJ: Prentice-Hall.
- Horowitz, M.J. (1986). *Stress Response Syndromes*. Northvale, NJ: Jason Aronson, Inc.
- Johnston, V. S., Miller, D. R., & Burleson, M. H. (1986). Multiple P3s to emotional stimuli and their theoretical significance. *Psychophysiology*, 23:684-694.
- Kagan, J. (1984). *The nature of the child*. New York: Basic Books.
- Kagan, J. (1994). *Galen's prophecy: Temperament in human nature*. New York: Basic Books.
- Kandel, E.R., Schwartz, J.H., & Jessell, T.M. (1991). *Principles of Neural Science*. New York, NY: Elsevier.
- Kardiner A. (1941). *The traumatic neuroses of war*. New York: Hoeber.
- Kaspi, S. P., McNally R. J., Amir, N. (1995). Cognitive processing of emotional information in post-traumatic stress disorder. *Cognitive Therapy and Research*, 19:433-444.
- Keane, T.M., and Kaloupek, D.G. (1982). Imaginal flooding in the treatment of a posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 50:138-140.
- Keane, T.M., Fairbank, J.A., Caddell, J.M., & Zimering, R.T. (1985). A behavioral approach to assessing and treating post-traumatic stress disorder in Vietnam veterans. In C.R. Figley (Ed.), *Trauma and its wake*. New York: Bruner Mazel, pp. 257-94.

- Keane, T.M., Fairbank, J.A., Cadell, J.M., Zimering, R.T., Taylor, K.L., Mora, C.A. (1989). Clinical evaluation of a measure to assess combat exposure: A psychological assessment. *Journal of Consulting and Clinical Psychology*, 1:53-55.
- Keane, T.M. (1993). Symptomatology in Vietnam Veterans with Post-traumatic Stress Disorder. In J.R.T. Davidson and E.B. Foa (Eds.), *Posttraumatic Stress Disorder: DSM-IV and Beyond*. Washington: American Psychiatric Press, pp. 99-112.
- Kendall-Tackett, K.A., Williams, L.M., Finkelhor, D. (1993). Impact of sexual abuse on children: A review and synthesis of recent empirical studies. *Psychological Bulletin*, 113:164-180.
- Kilpatrick, D.G., Veronen, L.J., & Best, C.L. (1985). Factors predicting psychological distress among rape victims. In C.R. Figley (Ed.), *Trauma and Its Wake*. New York: Brunner/Mazel.
- Kinard, E.M. (1995). Mother and teacher assessments of behavior problems in abused children. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34:1043-1053.
- Kindlon, D.J., Tremblay, R.E., Mezzacappa, E. (1995). Longitudinal patterns of heart rate and fighting behavior in 9 through 12-year-old boys. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34:371-377.
- Kiser, L.J., Heston, J., Millsap, P.A., Pruitt, D.B. (1991). Physical and sexual abuse in childhood: Relationship with post-traumatic stress disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 30(5):776-783.
- Kolb, L. (1984). The posttraumatic stress disorders of combat: A subgroup with a conditioned emotional response. *Military Medicine*, 149:237-243.
- Kolb L. (1987). A Neuropsychological Hypothesis Explaining Posttraumatic Stress Disorders. *American Journal of Psychiatry*, 144:989-995.
- Kostandov, E., & Arzumanov, Y. (1977). Averaged cortical evoked potentials to recognized and non-recognized verbal stimuli. *Acta Neurobiologiae Experimentalis*, 37:311-324.
- Kosten, T.R., Mason, J.W., Giller, E.L., Ostroff, R.B., & Harkness, L. (1987). Sustained urinary norepinephrine and epinephrine elevation in PTSD. *Psychoneuroendocrinology*, 12(1):13-20.
- Krystal, J., Kosten, T., Southwick, S., Mason, J.W., Perry, B. and Giller, E. (1989). Neurobiological Aspects of PTSD: Review of Clinical and Preclinical Studies. *Behavior Therapy*, 20:177-198.
- Kulka, R.A.; Schlenger, W.E.; Fairbank, J.A.; Hough, R.L.; Jordan, B.K.; Marmar, C.R.; Weiss, D.S. (1988). Contractual report of findings from the National Vietnam Veterans Readjustment Study: Executive Summary. Research Triangle Park, NC, Research Triangle Institute.
- Lang, P.J. (1984). *Emotions, cognition and behavior*. New York: Cambridge University Press.
- Lang, P.J. (1985). The cognitive psychophysiology of emotion: Fear and anxiety. In A.H. Tuma & J.D. Maser (Eds.), *Anxiety and the Anxiety Disorders*. Hillsdale, NJ: Erlbaum, pp. 131-70.
- Leichtman, M.D., Ceci, S.J., & Ornstein, P.S. (1992). The influence of affect on memory: Mechanism and development. In S. Christianson (Ed.), *The handbook of emotion and memory: Research and theory*. Hillsdale, NJ: Erlbaum, pp. 181-99.
- Lewis, D.L., and Hare, N. (1978). Review of the theoretical rationale and empirical support for the extinction approach of impulsive (flooding) therapy. In M. Hensen, R. Eisler, P. Miller (eds.), *Progress in Behavior Modification*. New York: Academic Press.
- Litz, B.T., & Keane, T.M. (1989). Information processing in anxiety disorders: Application to the understanding of post-traumatic stress disorder. *Clinical Psychology Review*, 9:243-257.
- Livingston, R., Lawson, L., Jones, J.G. (1993). Predictors of self-reported psychopathology in children abused repeatedly by a parent. *Journal of the American Academy of Child and Adolescent Psychiatry*, 32:948-953.
- MacLeod, C. (1990). Mood disorders and cognition. In M.W. Eysenck (Ed.), *Cognitive psychology: An international review*. New York: Wiley, pp. 9-56.
- MacLeod, C.M. (1991). Half a century of research on the Stroop effect: An integrative review. *Psychological Bulletin*, 109:163-203.
- Mack, R.L., Davenport, O.L., & Dykman, R.A. (1961). Cardiovascular conditioning in dogs. *American Journal of Physiology*, 201:437-439.
- Malloy, P.F., Fairbank, J.A., Keane, T.M. (1983). Validation of a multimethod assessment of post-traumatic stress disorders in Vietnam veterans. *Journal of Consulting Clinical Psychology*, 51:488-494.
- Mason, J., Giller, E., Kosten, T., Ostroff, R. and Podd, L. (1986). Urinary Free-Cortisol Levels in Posttraumatic Stress Disorder Patients. *Journal of Nervous and Mental Disease*, 174:145-149.
- McCarthy, G., & Donchin, E. (1981). A metric for thought: A comparison of P300 latency and reaction time. *Science*, 211:77-80.

- McFarlane, A.C. (1989). The aetiology of post-traumatic morbidity: Predisposing, precipitating and perpetuating factors. *British Journal of Psychiatry*, 154:221-228.
- McFarlane, A.C., Papay, P. (1992). Multiple diagnoses in posttraumatic stress disorder in the victims of a natural disaster. *Journal of Nervous and Mental Disease*, 180:498-504.
- McFarlane, A.C., Weber, D.L., Clark, R.C. (1993). Abnormal stimulus processing in posttraumatic stress disorder. *Biological Psychiatry*, 34:311-320.
- McGowan, W.T. III and Appel, J.B. (1977). An APL program for calculating signal detection indices. *Behavior Research Methods & Instrumentation*, 9:517.
- McLeer, S.V., Deblinger, E., Atkins, M.S., Foa, E.B., Ralphe, D.L. (1988). Posttraumatic stress disorder in sexually abused children. *Journal of the American Academy of Child and Adolescent Psychiatry*, 27:650-654.
- McLeer, S.V., Deblinger, E., Henry, D., Orvaschel, H. (1992). Sexually abused children at high risk for post-traumatic stress disorder. *Journal of the American Academy of Child and Adolescent Psychiatry*, 31:875-879.
- McLeer, S.V., Callaghan, M., Henry, D., Wallen, J. (1994). Psychiatric disorders in sexually abused children. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33:313-319.
- McNally, R.J., Luedke, D.L., Besyner, J.K., Peterson, R.A., Bohm, K., & Lips, O. J. (1987). Sensitivity to stress-relevant stimuli in posttraumatic stress disorder. *Journal of Anxiety Disorders*, 1:105-116.
- McNally, R.J., Kaspi, S.P., Riemann, B.C., Zeitlin, S.B. (1990). Selective processing of threat cues in Posttraumatic Stress Disorder. *Journal of Abnormal Psychology*, 99(4):398-402.
- McNally, R.J., English, G.E., Lipke, H.J. (1993). Assessment of intrusive cognition in PTSD: Use of the modified Stroop paradigm. *Journal of Traumatic Stress*, 6(1):33-38.
- McNally, R.J. and Saigh, P.A. (1993). On the distinction between Traumatic Simple Phobia and Posttraumatic Stress Disorder. In J.R.T. Davidson. and E.B. Foa (Eds.), *Posttraumatic Stress Disorder: DSM-IV and Beyond*. Washington: American Psychiatric Press, pp. 207-212.
- McPherson, W.B., Newton, J.E.O., Ackerman, P., Oglesby, D.M., & Dykman, R.A. (1996). An event-related brain potential investigation of PTSD and PTSD symptoms in abused children. *Integrative Physiological and Behavioral Science* (this volume).
- Mennen, F.E. (1993). Evaluation of risk factors in childhood sexual abuse. *Journal of the American Academy of Child and Adolescent Psychiatry*, 32:934-939.
- Merry, S.N. & Andrews, L.K. (1994). Psychiatric status of sexually abused children 12 months after disclosure of abuse. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33:939-944.
- Metzger, L.J., Orr, S.P., McNally, R.J., and Pitman, R.K. (1996). Seeking the source of emotional stroop interference effects in PTSD: A study of P3s to traumatic words. *Integrative Physiological and Behavioral Sciences* (this volume).
- Metzger, L.J., Orr, S. P., Lasko, N.B., & Pitman, R.K. (in press). Event-related potentials to tone stimuli in combat-related post-traumatic stress disorder. *Biological Psychiatry*.
- Meuser, K.T., Yarnold, P.R., Foy, D.W. (1991). Statistical analysis for single-case designs: Evaluating outcome of imaginal exposure treatment of chronic PTSD. *Behavior Modification*, 15:134-155.
- Mogg, K. & Marden, B. (1990). Processing of emotional information in anxious subjects. *British Journal of Clinical Psychology*, 29:227-229.
- Mowrer, O.H. (1947). On the dual nature of learning: A reinterpretation of "conditioning" and "problem solving." *Harvard Education Review*, 17:102-148.
- Mowrer, O.H. (1960). *Learning Theory and the Symbolic Processes*. New York: John Wiley & Sons.
- Murphree, O.D. & Dykman, R.A. (1965). Litter patterns in the offspring of nervous and stable dogs. I: Behavioral tests. *Journal of Nervous and Mental Disease*, 141:321-332.
- Murphree, O.D., Dykman, R.A., & Peters, J.E. (1967). Operant conditioning of two strains of pointer dogs. *Psychophysiology*, 4:414-417.
- Naatanen, R.(1982). Processing negativity: An evoked-potential reflection of selective attention. *Psychological Bulletin*, 92:605-640.
- Newton, J.E.O. & Gantt, W.H. (1966). One-trial cardiac conditioning in dogs. *Conditional Reflex*, 1:18-35.
- Newton, J.E.O., Chapin, J.L., & Murphree, O.D. (1976). Correlations of normality and nervousness with cardiovascular functions in pointer dogs. *Pavlovian Journal*, 11.
- Ornitz, E.M. & Pynoos, R.S. (1989). Startle modulation in children with posttraumatic stress disorder. *American Journal of Psychiatry*, 146:866-870.

- Orr, S.P. (1990). Psychophysiological studies of post traumatic stress disorder. In E.L. Giller (Ed). *Biological Assessment and Treatment of Post Traumatic Stress Disorder*. Washington: American Psychiatric Press, Inc., pp. 75-90.
- Orvaschel, H., Puig-Antich, P., Chambers, W., Abrizzi, M. (1982). Retrospective assessment of prepubertal major depression with the Kiddie SADS. *Journal of the American Academy of Child and Adolescent Psychiatry*, 21:392-397.
- Paige, S.R., Reid, G.M., Allen, M.G., & Newton, J.E.O. (1990). Psychophysiological correlates of posttraumatic stress disorder in Vietnam veterans. *Biological Psychiatry*, 27:419-430.
- Paige, S.R., Fitzpatrick, D.F., Kline, J.P., Balogh, S.E., & Hendricks, S.E. (1994). Event-related potential amplitude/intensity slopes predict response to antidepressants. *Neuropsychobiology*, 30:197-201.
- Pallmeyer, T., Blanchard, E. and Kolb, L. (1986). The Psychophysiology of Combat-Induced Post-Traumatic Stress Disorder in Vietnam Veterans. *Behavior Research and Therapy*, 24:645-652.
- Pary, R., Tobias, C. and Lippman, S. (1987). Recognizing Shammed and Genuine Post-Traumatic Stress Disorder. *VA Practitioner* (July):37-43.
- Pavlov, I.P. (1927). *Conditioned Reflexes* (Trans. by G.V. Anrep.). London: Oxford University Press.
- Pavlov, I.P. (1928). *Lectures on Conditioned Reflexes* (Trans. by W.H. Gantt.). New York: International Publishers.
- Pelcovitz, D., Kaplan, S., Goldenberg, B., Mandel, F., Lehane, J., Guarrera, J. (1994). Post-traumatic stress disorder in physically abused adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33:305-312.
- Perry, B.D. (1994). Catecholamine function in posttraumatic stress disorders: Emerging concepts. In M. Murbery (Ed.), *Neurobiological Sequelae of Childhood Trauma*. Washington, DC: American Psychiatric Press, pp. 221-56.
- Pfefferbaum, A., Wenegrat, B.G., Ford, J.M., Walton, T., Roth, W.T., & Kopell, B.S. (1984). Clinical application of the P3 component of event-related potentials: II. Dementia, depression and schizophrenia. *Electroencephalography and Clinical Neurophysiology*, 59:104-124.
- Pitman, R.K. & Orr, S.P. (1986). Test of the conditioning model of neurosis: Differential aversive conditioning of angry and neutral facial expressions in anxiety disorder patients. *Journal of Abnormal Psychology*, 95:208-213.
- Pitman, R.K., Orr, S.P., Foa, D.F., DeJong, J.B., & Claiborn, J.M. (1987). Psychophysiological assessment of post-traumatic stress disorder in Vietnam combat veterans. *Archives of General Psychiatry*, 44:970-975.
- Pitman, R.K. (1988). Post-Traumatic Stress Disorder, Conditioning, and Network Theory. *Psychiatric Annals*, 18:182-189.
- Pitman, R.K. (1989). Post-Traumatic Stress Disorder, Hormones, and Memory. *Biological Psychiatry*, 26:221-223.
- Pitman, R.K. and Orr, S.P. (1990). Twenty-Four Hour Urinary Cortisol and Catecholamine Excretion in Combat-Related Posttraumatic Stress Disorder. *Biological Psychiatry*, 27:245-247.
- Pitman, R.K., Orr, S.P., Foa, D.F., Altman, B., de Jong, J.B., Herz, L.R. (1990). Psychophysiological responses to combat imagery of Vietnam veterans with posttraumatic stress disorder versus other anxiety disorders. *Journal of Abnormal Psychology*, 99(1):49-54.
- Pitman, R., Altman, B., Greenwald, E. (1991). Psychiatric complications during flooding therapy for posttraumatic stress disorder. *Journal of Clinical Psychiatry*, 52:17-20.
- Pitman, R.K. (1993). Biological Findings in Posttraumatic Stress Disorder: Implications for DSM-IV Classification. In J.R.T. Davidson and E.B. Foa (Eds.), *Posttraumatic Stress Disorder: DSM-IV and Beyond*. Washington: American Psychiatric Press, pp.173-190.
- Porges, S.W. (1995). Cardiac vagal tone: A physiological index of stress. *Neuroscience and Biobehavioral Reviews*, 19:225-233.
- Prescott, J., Connolly, J.F., & Gruzelier, J.H. (1984). The augmenting/reducing phenomenon in the auditory-evoked potential. *Biological Psychology*, 19:31-44.
- Prins, A.T.; Kaloupek, D.G.; Keane, T.M. (1995). Psychophysiological evidence of autonomic arousal and startle in traumatized adult populations. In M. Friedman, D. Charney, A. Deutch (Eds), *Neurobiological and clinical consequences of stress: From normal adaptation to PTSD*. Philadelphia: Lippincott-Raven Publishers, pp. 291-314.
- Pritchard, W.S. (1981). Psychophysiology of P300. *Psychological Bulletin*, 89:506-540.
- Pritchard, W.S. (1986). Cognitive event-related potentials correlates of schizophrenia. *Psychological Bulletin*, 100:43-66.

- Pynoos, R.S.; Eth, S. (1985). Children traumatized by witnessing acts of personal violence: Homicide, rape, or suicide behavior. In S. Eth & R.S. Pynoos (Eds.), *Post-traumatic stress disorder in children*. Washington, DC: American Psychiatric Press, pp. 19-43.
- Pynoos, R.S., Frederick, C., Nader, K., Arroyo, W., Steinberg, A., Eth, S., Nunez, F., & Fairbanks, L. (1987). Life threat and posttraumatic stress in school-age children. *Archives of General Psychiatry*, 44:1057-1063.
- Raine, A., Venables, P.H., Williams, M. (1990). Relationships between central and autonomic measures of arousal at age 15 years and criminality at age 24 years. *Archives of General Psychiatry*, 47:1003-1007.
- Ray, C. (1979). Examination stress and performance on a color-word interference test. *Perceptual and Motor Skills*, 49:400-402.
- Reich, W. & Welner, Z. (1988). *DICA-R-C (DSM-III-R Version), Revised Version 5-R*. St. Louis, MO: Washington University.
- Rothbaum, B.O. and Foa, E.B. (1993). Subtypes of posttraumatic stress disorder and duration of symptoms. In J.R.T. Davidson and E.B. Foa (Eds.), *Posttraumatic Stress Disorder: DSM-IV and Beyond*. Washington: American Psychiatric Press, pp. 23-36.
- Rutschmann J, Cornblatt B, Erlenmeyer-Kimling L (1977). Sustained Attention in Children at Risk for Schizophrenia. Report on a Continuous Performance Test. *Archives of General Psychiatry*, 34:571-75.
- Rutter, M. & Graham, P. (1968). The reliability and validity of the psychiatric assessment of the child: I. Interview with the child. *British Journal of Psychiatry*, 114:563-579.
- Scerbo, A.S. & Kolko, D.J. (1995). Child physical abuse and aggression: Preliminary findings on the role of internalizing problems. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34:1060-1066.
- Scherg, M. and Von Cramon, D. (1985). Two bilateral sources of the late AEP as identified by a spatio-temporal dipole model. *Electroencephalography and Clinical Neurophysiology*, 62:32-44.
- Seligman, M.E. (1970). On the generality of the laws of learning. *Psychological Review*, 77:406-418.
- Seligman, M.E. (1971). Phobias and preparedness. *Behavior Therapy*, 2:307-320.
- Shalev, A.Y., Orr, S.P., Peri, T., Schreiber, S., & Pitman, R.K. (1992). Physiologic responses to loud tones in Israeli patients with posttraumatic stress disorder. *Archives of General Psychiatry*, 49:870-875.
- Solomon, S.D., Gerrity, E.T., Muff, A.M. (1992). Efficacy of treatment for posttraumatic stress disorder. *Journal of the American Medical Association*, 268:633-638.
- Southwick, S.M., Krystal, J.H., Morgan, C.A., Johnson, D., Nagy, L.M., Nicolaou, A., Heninger, G.R., & Charney, D.S. (1993). Abnormal noradrenergic function in posttraumatic stress disorder. *Archives of General Psychiatry*, 50:266-274.
- Sparr, L. and Pankrantz, L.D. (1983). Factitious post-traumatic stress disorder. *American Journal of Psychiatry*, 140:1016-1019.
- Spielberger, C. D., Gorsuch, R. L., Lushene, R. E., Vagg, P.R., & Jacobs, G. A. (1983): *Manual for the State-Trait Anxiety Inventory* (Self-Evaluation Questionnaire). Palo Alto: Consulting Psychologists Press.
- Spitzer, R.L., Williams, J.B., Gibbon, M. (1987). *Structured Clinical Interview for DSM-III-R*. New York: New York State Psychiatric Institute.
- Spitzer, R.L., Williams, J.B., Gibbon, M., First, M.B. (1990) Structured Clinical Interview for the DSM-III-R-Patient Edition (SCID-P), *Biometrics Research*.
- Spitzer, R. L., Williams, J. B., Gibbon, M., & First, M. B. (1992): The Structured Clinical Interview for DSM-III-R (SCID): I. History, rationale, and description. *Archives of General Psychiatry*, 49:624-629.
- Stampfl, T.G. & Levis, D.J. (1967). Essentials of implosive therapy: A learning-theory-based psychodynamic behavioral therapy. *Journal of Abnormal Psychology*, 72:157-163.
- Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, 18:643-662.
- Sutker, P.B., Winstead, D.K., Galina, Z.H., Allain, A.N. (1991). Cognitive deficits and psychopathology among former prisoners of war and combat veterans of the Korean conflict. *American Journal of Psychiatry*, 148:67-72.
- Swets, J.A., Tanner, W.P. Jr, Birdsall, T.G. (1961). Decision processes in perception. *Psychological Review*, 68:301-340.
- Terr, L.C. (1991). Childhood traumas: An outline and overview. *American Journal of Psychiatry*, 148:10-20.
- Thrasher, S. M., Dalgleish, T., & Yule, W. (1994). Information processing in posttraumatic stress disorder. *Behaviour Research and Therapy*, 32:247-254.
- Trickett, P.K. & Putnam, F.W. (1993). Impact of child sexual abuse on females: Toward a developmental, psychobiological integration. *Psychological Science*, 4:81-87.

- True, W.R., Rice, J., Eisen, S.A., Heath, A.C., Goldberg, J., Lyons, M.J., & Nowak, J. (1993). A twin study of genetic and environmental contributions to liability for posttraumatic stress symptoms. *Archives of General Psychiatry*, 50:257-264.
- van der Kolk, B., Greenberg, M., Boyd, H. and Krystal, J. (1985). Inescapable Shock, Neurotransmitters and Addiction to Trauma: Toward a Psychobiology of Post Traumatic Stress Disorder. *Biological Psychiatry*, 20:314-325.
- Verleger, R. (1988). From Epistemology to P3-ology. *Behavior and Brain Science*, 11(3):399-427.
- Vrana, S. R., Roodman, A., & Beckham, J. C. (1995). Selective processing of trauma-relevant words in posttraumatic stress disorder. *Journal of Anxiety Disorders*, 9:515-530.
- Warren, L. R., & Marsh, G. R., (1979). Changes in event related potentials during processing of Stroop stimuli. *International Journal of Neuroscience*, 9:217-223.
- Wells, A. & Matthews, G. (1994). *Attention and emotion: A clinical perspective*. Hillsdale, NJ: Erlbaum.
- Williams, J.M.G., Watts, F.N., MacLeod, C., & Mathews, A. (1988). *Cognitive psychology and emotional disorders*. New York: Wiley.
- Williamson, S. Harpur, T., & Hare, R. D. (1991). Abnormal Processing of Affective Words by Psychopaths. *Psychophysiology*, 28:260-273.
- Woodward, S.H., Drescher, K.D., Murphy, R.T., Ruzek, J.I., Foy, D.W., Arsenault, N.J., and Gusman, F.D. (1996). Heart rate during group flooding therapy for PTSD. *Integrative Physiological and Behavioral Sciences* (this volume).
- Yalom, I.D. (1985). *The theory and practice of group psychotherapy*. (3rd edition). New York: Basic Books.
- Yehuda, R., Giller, E.L., Southwick, S., Lowy, M.T. and Mason, J.W. (1991). Hypothalamic pituitary adrenal dysfunction in posttraumatic stress disorder. *Biological Psychiatry*, 30:1031-1048.
- Yehuda, R., Southwick, S.M., Ma, X., Giller, E.L., & Mason, J.W. (1992). Urinary catecholamine excretion and severity of symptoms in PTSD. *Journal of Nervous and Mental Disorders*, 180:321-325.
- Yehuda, R., Kahana, B., Binder-Brynes, K., Southwick, S.M., Mason, J.W., & Giller, E.L. (1995a). Low urinary cortisol excretion in holocaust survivors with posttraumatic stress disorder. *American Journal of Psychiatry*, 152:982-985.
- Yehuda, R., Keefe, R.S.E., Harvey, P.D., Levengood, R.A., Gerber, D.K., Geni, J., and Siever, L.J. (1995b). Learning and memory in combat veterans with posttraumatic stress disorder. *American Journal of Psychiatry*, 152:137-139.
- Yehuda, R. & McFarlane, A.C., (1995). Conflict between current knowledge about posttraumatic stress disorder and its original conceptual basis. *American Journal of Psychiatry*, 152:1705-1713.
- Zaidi, L.Y. & Foy, D.W. (1994). Childhood abuse experiences and combat-related PTSD. *Journal of Traumatic Stress*, 7:33-42.
- Zeitlin, S.B., McNally, R.J. (1991). Implicit and explicit memory bias for threat in post-traumatic stress disorder. *Behavior Research and Theory*, 29(5):451-457.